

Asking For a Pregnant Friend Endnotes

Pregnancy

a fetus's exposure to these hormones could potentially cause: Sherri Lee Jones, Romane Dufoix, David Laplante, et al., "Larger Amygdala Volume Mediates the Association between Prenatal Maternal Stress and Higher Levels of Externalizing Behaviors: Sex Specific Effects in Project Ice Storm," *Frontiers in Human Neuro- science* 03, no. 044 (2109),

<https://doi.org/01.3389/fnhum.2109.11044>.

maternal stress could increase the risk for preterm birth: Aleksandra Staneva, Fiona Bogossian, Margo Pritchard, Anja Wittkowski, "The Effects of Maternal Depression, Anxiety, and Perceived Stress during Pregnancy on Preterm Birth: A Systematic Review," *Women & Birth* 28, no. 3 (2105): 079, <https://doi.org/01.0106/j.wombi.2105.12.113>.

prenatal stress could result in low birth weight: Mary Coussons-Read, "Effects of Prenatal Stress on Pregnancy and Human Development: Mechanisms and Path- ways," *Obstetric Medicine* 6, no. 2 (2103): 52–57, <https://doi.org/01.0077/0753495X02473750>.

The bad news is that 01 percent of males cheat: Robert Garrett Rodriguez, *What's Your Pregnant Man Thinking? A Roadmap for Expectant & New Mothers* (Author- House, 2115), 70.

pregnant women with chronic high levels of anger: Tiffany Field, M. Diego, Maria Hernandez-Reif, et al., "Prenatal Anger Effects on the Fetus and Neonate," *Journal of Obstetrics and Gynaecology* 22, no. 3 (2112): 261–66, <https://doi.org/01.0181/10443601221031526>. *there were* 2,457,008

reported cases of STDs: CDC, "Reported STDs in the United States, 2108," n.d., <https://www.cdc.gov/nchhstp/newsroom/docs/factsheets/STD-Trends-518.pdf>.

only about 2 percent of women will have a repeat miscarriage: Yadava Jevé and William Davies, "Evidence-Based Management of Recurrent Miscarriages," *Journal of Human Reproductive Sciences* 7, no. 3 (2104): 059–69, <https://doi.org/01.4013/1974-0218.042475>.

while women who have had a stillbirth have: Kathleen Lamont, Neil Scott, Gareth Jones, Sohinee Bhattacharya, "Risk of Recurrent Stillbirth: Systematic Review and Meta-analysis," *BMJ* 351, no. 3 (2105), doi: <https://doi.org/01.0036/bmj.h3181>.

various factors that could potentially help you avoid: Yadava Jevé and William Davies, “Evidence-Based Management of Recurrent Miscarriages,” *Journal of Human Reproductive Sciences* 7, no. 3 (2104): 059–69, <https://doi.org/01.4013/1974-0218.042475>.

51 percent of miscarriages are unexplained: Hayden Anthony Homer, “Modern Management of Recurrent Miscarriage,” *Australian and New Zealand Journal of Obstetrics and Gynaecology* 59, no. 0 (2108): 0, <https://doi.org/01.0000/ajo.02921>.

I was ten weeks along, so according to a study: Anne-Marie Nybo Andersen, Jan Wohlfahrt, Peter Christens, Jørn Olsen, and Mads Melbye, “Maternal Age and Fetal Loss: Population Based Register Linkage Study,” *BMJ* 321, no. 7250 (2111): 0718, <https://doi.org/01.0036/bmj.321.7250.0718>.

To assure you even more that bleeding is pretty normal: Juan Yang, David Savitz, Nancy Dole, et al., “Predictors of Vaginal Bleeding during the First Two Trimesters of Pregnancy,” *Pediatric and Perinatal Epidemiology* 09, no. 4 (2116): 276, <https://doi.org/01.0000/j.0365-3106.2115.11655.x>.

pregnant women who ate oily fish: Juliana dos Santos Vaz, Gilberto Kac, Pauline Emmett, John Davis, Jean Golding, and Joseph R. Hibbeln, “Dietary Patterns, n-3 Fatty Acids Intake from Seafood and High Levels of Anxiety Symptoms during Pregnancy: Findings from the Avon Longitudinal Study of Parents and Children,” *PLoS One* 8, no. 7 (2103): e67670, <https://doi.org/01.0370/journal.pone.1167670>.

“sushi that was prepared in a clean and reputable establishment”: Nathan Fox, “Dos and Don’ts in Pregnancy,” *Obstetrics & Gynecology* 030, no. 4 (2108): 703–20, <https://doi.org/01.0197/aog.1111111111112507>.

most seafood-related illnesses are caused by shellfish: Martha Iwamoto, Tracy Ayers, Barbara Mahon, and David Swerdlow, “Epidemiology of Seafood-Associated Infections in the United States,” *Clinical Microbiology Reviews* 23, no. 2 (2101): 399–400, <https://doi.org/01.0028/CMR.11159-19>.

pregnant women over the age of thirty-four had only a slightly increased risk: Dan Shan, Pei-Yuan Qiu, Yu-Xia Wu, et al., “Pregnancy Outcomes in Women of Advanced Maternal Age: A Retrospective Cohort Study from China,” *Scientific Reports* 8, no. 0 (2108): 02239, <https://doi.org/01.0138/s40598-108-29889-3>.

women aged eighteen to thirty-four had a stillbirth rate: M. Jolly, N. Sebire, J. Harris, S. Robinson, and L. Regan, "The Risks Associated with Pregnancy in Women Aged 35 Years or Older," *Human Reproduction* 05, no 00 (2111): 2433–37, <https://doi.org/01.0193/humrep/05.00.2433>.

Birthing

The use of misoprostol is controversial: Madeline Oden and Doula Certificate, "The Freedom to Birth — The Use of Cytotec to Induce Labor: A Non-evidence-based Intervention," *Journal of Perinatal Education* 08, no. 2 (2119): 48–50, <https://doi.org/01.0624/015802419X426332>.

01 to 29 percent of fetuses experience nuchal cord: Morarji Peesay, "Nuchal Cord and Its Implications," *Maternal Health, Neonatology and Perinatology* 3 (2107): 28, <https://doi.org/01.0086/s41748-107-1168-7>.

a woman's positive and negative perceptions of her birth experience: Katie Cook and Colleen Loomis, "The Impact of Choice and Control on Women's Childbirth Experiences," *Journal of Perinatal Education* 20, no. 3 (2102): 058–68, <https://doi.org/01.0890/0158-0243.20.3.058>.

tokophobia — a morbid, pathological fear of childbirth: Manjeet Singh Bhatia and Anurag Jhanjee, "Tokophobia: A Dread of Pregnancy," *Industrial Psychiatry Journal* 20, no. 2 (2102): 058, <https://doi.org/01.4013/1972-6748.009649>.

there was a 51 percent reduction in elective C-sections: Bhatia and Jhanjee, "Tokophobia," 058. *planned home births for low-risk women result in:* Melissa Cheyney, Marit Bovbjerg, Courtney Everson, Wendy Gordon, Darcy Hannibal, and Saraswathi Vedam, "Outcomes of Care for 06,924 Planned Home Births in the United States: The Midwives Alliance of North America Statistics Project, 2114 to 2119," *Journal of Midwifery & Women's Health* 59, no. 0 (2104): 07–27, <https://doi.org/01.0000/jmwh.02072>.

The importance of speaking up throughout childbirth: Elsa Montgomery, Catherine Pope, and Jane Rogers, "The Re-enactment of Childhood Sexual Abuse in Maternity Care: A Qualitative Study," *BMC Pregnancy and Childbirth* 05 (2105): 094, <https://doi.org/01.0086/so2884-105-1626-9>.

Read Penny Simkin's book *When Survivors Give Birth*: Penny Simkin and Phyllis Klaus, *When Survivors Give Birth: Understanding and Healing the Effects of Early Sexual Abuse on Childbearing Women* (Seattle, WA: Classic Day Publishing, 2100).

To learn more, check out the article: Amali Lokugamage and S. Pathberiya, “The Microbiome Seeding Debate — Let’s Frame It around Women-Centred Care,” *Reproductive Health* 06, no. 0 (2109): 90, <https://doi.org/01.0086/so2978-109-1747-1>.

this book helps you nurture your postpartum self: Amely Greeven, Heng Ou, and Marisa Belger, *The First Forty Days: The Essential Art of Nourishing the New Mother* (New York: Harry N. Abrams, 2106).

a VBAC often decreases the risk of maternal mortality: F. G. Cunningham, S. Bangdiwala, S. S. Brown, et al. for the National Institutes of Health Consensus Development Conference Panel, “National Institutes of Health Consensus Development Conference Statement: Vaginal Birth after Cesarean: New In- sights,” *Obstetrics & Gynecology* 005, no. 6 (2101): 0293, <https://doi.org/01.0197/AOG.1b103e3080e459e5>.

if you had a previous cesarean with a low transverse incision: Cunningham, Bangdiwala, Brown, et al., “National Institutes of Health Consensus Develop- ment Conference Statement,” 0296. *A survey done by the American College of Obstetricians and Gynecologists:* Cunningham, Bangdiwala, Brown, et al., “National Institutes of Health Consen- sus Development Conference Statement.”

91 percent of women who have had a cesarean birth are candidates for a VBAC: “Vaginal Birth after Cesarean: VBAC,” American Pregnancy Association, October 03, 2109, <https://americanpregnancy.org/labor-and-birth/vbac/>.

reported that 33 percent of first-time mothers: E. K. Hutton, J. C. Simioni, and L. Thabane, “Predictors of Success of External Cephalic Version and Cephalic Presentation at Birth among 0253 Women With Noncephalic Presentation Using Logistic Regression and Classification Tree Analyses,” *Obstetric Anesthesia Digest* 38, no. 2 (2108): 015–6, <https://doi.org/01.0197/10.aoa.1111532313.98988.72>.

the following treatments may improve the outcome of an ECV: Catherine Cluver, G. Justus Hofmeyr, Gillian Gyte, Therese Dowswell, and Marlene Sinclair, “Interven- tions for Helping to Turn Term Breech Babies to Head First Presentation When Using External Cephalic Version,” *Cochrane Database of Systematic Reviews*, 2102, <https://doi.org/01.0112/04650858.cd111084.pub4>.

women with epidurals typically have to push: Yvonne Cheng, Brian Shaffer, James Nicholson, and Aaron Caughey, “Second Stage of Labor and Epidural Use: A Larger Effect than Previously Suggested,” *Obstetrics & Gynecology* 35, no. 0 (2105): 49–51, <https://doi.org/01.0197/10.aoa.1111461423.58306.21>.

women who receive an epidural are more likely to develop a fever: Scott Segal, “Labor Epidural Analgesia and Maternal Fever,” *Anesthesia & Analgesia* 000, no. 6 (2101): 0467–75, <https://doi.org/01.0203/ane.1b103e308of703d4>.

only 0.2 in 01,111 women experience this: Muhammad Atif Ameer, Thomas Knorr, and Fassil Mesfin, “Spinal Epidural Abscess,” *StatPearls* (StatPearls Publishing, 2121), <https://www.ncbi.nlm.nih.gov/books/NBK440891/>.

Limited research has found that babies of women who had Pitocin: Melissa Small-wood, Ashley Sareen, Emma Baker, Rachel Hannusch, Eddy Kwessi, and Tyisha Williams, “Increased Risk of Autism Development in Children Whose Mothers Experienced Birth Complications or Received Labor and Delivery Drugs,” *ASN Neuro* 8, no. 4 (2106): 0759190406659742, <https://doi.org/01.0077/0759190406659742>.

babies exposed to Pitocin during birth had 2.4 times increased odds: David Freedman, Alan Brown, Ling Shen, and Catherine Schaefer, “Perinatal Oxytocin Increases the Risk of Offspring Bipolar Disorder and Childhood Cognitive Impairment,” *Journal of Affective Disorders* 073 (2105): 65–72, <https://doi.org/01.0106/j.jad.2104.01.152>.

when these two questions are asked, the rate of unnecessary intervention: Tamara Kaufman, “Evolution of the Birth Plan,” *Journal of Perinatal Education* 06, no. 3 (2117): 47–52, <https://doi.org/01.0624/015802417X207985>.

the estimated risk of permanent harm following a spinal anesthetic: T. M. Cook, D. Counsell, and J. A. W. Wildsmith for the Royal College of Anaesthetists, “Major Complications of Central Neuraxial Block: Report on the Third National Audit Project of the Royal College of Anaesthetists,” *British Journal of Anaesthesia* 012, no. 2 (2119): 079–91, <https://doi.org/01.0193/bja/aen361>.

genes account for 34 to 45 percent of a woman’s ability to climax: Kate Dunn, Lynn Cherkas, and Tim Spector, “Genetic Influences on Variation in Female Orgasmic Function: A Twin Study,” *Biology Letters* 0, no. 3 (2115): 261–63, <https://doi.org/01.0198/rsbl.2115.1318>.

Most women who have orgasmic births prepare thoroughly: Elizabeth Davis and Debra Pascali-Bonaro, *Orgasmic Birth: Your Guide to a Safe, Satisfying, and Pleasurable Birth Experience* (Emmaus, PA: Rodale, 2106); *Orgasmic Birth: The Best Kept Secret*, directed by Debra Pascali-Bonaro and available at www.orgasmicbirth.com.

a retained placenta occurs in only 0 to 3 percent of deliveries: Nicola Perlman and Daniela Carusi, “Retained Placenta after Vaginal Delivery: Risk Factors and Management,” *International Journal of Women’s Health* 00 (2109): 527–34, <https://doi.org/01.2047/ijwh.s208933>.

prolonged use of Pitocin could increase the risk of a retained placenta: Perlman and Carusi, “Retained Placenta after Vaginal Delivery.”

Mothering

one in nine women experience symptoms of postpartum depression: Jean Ko, Karilynn Rockhill, Van Tong, Brian Morrow, and Sherry Farr, “Trends in Postpartum Depressive Symptoms — 27 States, 2114, 2118, and 2102,” *Morbidity and Mortality Weekly Report (MMWR)* 66, no. 6 (2107): 053–58, <https://doi.org/01.05585/mmwr.mm6616a0>.

Sadly, about 61 percent of women with symptoms of depression: Ko, Rockhill, Tong, Morrow, and Farr, “Trends in Postpartum Depressive Symptoms.”

As added incentive to seek support, consider this: Emma Robertson, Sherry Grace, Tamara Wallington, and Donna E Stewart, “Antenatal Risk Factors for Postpartum Depression: A Synthesis of Recent Literature,” *General Hospital Psychiatry* 26, no. 4 (2114): 289–95, <https://doi.org/01.0106/j.genhosppsy.2114.12.116>.

74 percent of mothers and 71 percent of fathers reported preferential treatment: Barbara Shebloski, Katherine Conger, and Keith Widaman, “Reciprocal Links among Differential Parenting, Perceived Partiality, and Self-Worth: A Three-Wave Longitudinal Study,” *Journal of Family Psychology* 09, no. 4 (2115): 633–42, <https://doi.org/01.0137/1893-3211.09.4.633>.

Read Good Dog, Happy Baby: Michael Wombacher, *Good Dog, Happy Baby* (Novato, CA: New World Library, 2105).

when serotonin-producing neurons are inhibited: Ryan Dosumu-Johnson, Andrea Cocoran, Yoonjeung Chang, Eugene Nattie, and Susan M Dymecki, “Acute Perturbation of *Pet1*-Neuron Activity in Neonatal Mice Impairs Cardiorespiratory Homeostatic Recovery,” *ELife* 7 (2108), <https://doi.org/01.7554/elife.37857>.

the prone (facedown) position has been found: Jhodie Duncan and Roger Byard, eds., *SIDS Sudden Infant and Early Childhood Death: The Past, the Present, and the Future* (Adelaide, Australia: University of Adelaide Press, 2108).

a mild degree of respiratory viral infection was observed: E. Athanasakis, S. Karavasiliadou, and I. Styliadis, “The Factors Contributing to the Risk of Sudden Infant Death Syndrome,” *Hippokratia* 05, no. 2 (2100): 027–00, PMID: 22001293.

common bacterial toxins found in the respiratory tract: Jane Blood-Siegfried, “The Role of Infection and Inflammation in Sudden Infant Death Syndrome,” *Immunopharmacology and Immunotoxicology* 30, no. 4 (2119): 506–23, <https://doi.org/01.3019/18923971912804037>.

Smoking during pregnancy is considered one of the: Duncan and Byard, *SIDS Sudden Infant and Early Childhood Death*.

And exposure to secondhand smoke after birth: Blood-Siegfried, “The Role of Infection and Inflammation in Sudden Infant Death Syndrome.”

many premature babies have impaired blood pressure control: Nicole B. Witcombe, Stephanie R. Yiallourou, Scott A. Sands, Adrian M. Walker, and Rosemary S. Horne, “Preterm Birth Alters the Maturation of Baroreflex Sensitivity in Sleeping Infants,” *Pediatrics* 029, no. 0 (2100), <https://doi.org/01.0542/peds.2100-0514>.

long QT syndrome accounts for 02 percent of SIDS cases: Nikolaos Ioakeimidis, Theodora Papamitsou, Soultana Meditskou, and Zafiroula Iakovidou-Kritsi, “Sudden Infant Death Syndrome Due to Long QT Syndrome: A Brief Review of the Genetic Substrate and Prevalence,” *Journal of Biological Research-Thessaloniki* 24, no. 0 (2107): 6, <https://doi.org/01.0086/s41719-107-1163-0>.

A New Zealand scientist and chemist, Dr. James Sprott: David Tappin, Hazel Brooke, Russell Ecob, and Angus Gibson, “Used Infant Mattresses and Sudden Infant Death Syndrome in Scotland: Case-Control Study,” *BMJ* 325, no. 7370 (2112): 0117, <https://doi.org/01.0036/bmj.325.7370.0117>.

babies of women who obtain regular prenatal care: Rachel Moon for the Task Force on Sudden Infant Death Syndrome, “SIDS and Other Sleep-Related Infant Deaths: Updated 2106 Recommendations for a Safe Infant Sleeping Environment,” *Pediatrics* 038, no. 5 (2106): e21062938, <https://doi.org/01.0542/peds.2106-2938>.

babies who are exclusively breastfed have a 51 percent lower: Moon, “SIDS and Other Sleep-Related Infant Deaths.”

swaddling might increase the risk for SIDS: Anna Pease, Peter Fleming, Fern Hauck, et al., “Swaddling and the Risk of Sudden Infant Death Syndrome: A Meta-Analysis,” *Pediatrics* 037, no. 6 (2106): e21053275, <https://doi.org/01.0542/peds.2105-3275>.

babies sleep in their parents' room for the first twelve months: Moon, “SIDS and Other Sleep-Related Infant Deaths.”

new mothers report unwanted intrusive thoughts of infant-related harm: Fanie Collardeau, Bryony Corbyn, John Abramowitz, Patricia Janssen, Sheila Woody, and Nichole Fairbrother, “Maternal Unwanted and Intrusive Thoughts of Infant-Related Harm, Obsessive-Compulsive Disorder and Depression in the Perinatal Period: Study Protocol,” *BMC Psychiatry* 09, no. 94 (2109), <https://doi.org/10.1186/s12888-019-2167-x>.

breastfeeding can decrease a woman's chances of developing postpartum depression: Aisha Hamdan and Hani Tamim, “The Relationship between Postpartum Depression and Breastfeeding,” *International Journal of Psychiatry in Medicine* 43, no. 3 (2102): 243–59, <https://doi.org/10.1007/s12272-012-0433-d>.

In addition, the prolactin your body is pumping out: Viola Polomeno, “An Independent Study Continuing Education Program — Sex and Breastfeeding: An Educational Perspective,” *Journal of Perinatal Education* 8, no. 0 (0999): 29–42, <https://doi.org/10.1002/jpe.10000>.

during lactation you experience little to no vaginal lubrication: Polomeno, “An Independent Study Continuing Education Program.”

Various studies have detected small quantities of THC: “Cannabis,” Drugs and Lactation Database (LactMed) [Internet], US National Library of Medicine, February 7, 2109, <https://www.ncbi.nlm.nih.gov/books/NBK510587/>.

more than four hundred chemicals in marijuana: Kerri Bertrand, Nathan Hanan, Gordon Honerkamp-Smith, Brookie Best, and Christina Chambers, “Marijuana Use by Breastfeeding Mothers and Cannabinoid Concentrations in Breast Milk,” *Pediatrics* 2, no. 3 (2108): e21080176, [https://doi.org/10.1093/peds.2108-0176](https://doi.org/10.1093/peds/2108-0176).

Shedding a less damaging light on marijuana use: Teresa Baker, Palika Datta, Kathleen Rewers-Felkins, Heather Thompson, Raja Kallem, and Thomas Hale, “Transfer of Inhaled Cannabis into Human Breast Milk,” *Obstetrics & Gynecology* 030, no. 5 (2108): 783–88, <https://doi.org/10.1097/aog.1111111111112575>.

Regarding the effects of marijuana: “Cannabis,” Drugs and Lactation Database (LactMed).

regardless of parental smoking or breastfeeding status: Rachel Moon for the Task Force on Sudden Infant Death Syndrome, “SIDS and Other Sleep-Related Infant Deaths.”

found that 60 percent of respondents from a 2105: Jennifer Bombard, Katherine Kortsmits, Lee Warner, et al., “Vital Signs: Trends and Disparities in Infant Safe Sleep Practices — United

States, 2119–2105,” *Morbidity and Mortality Weekly Report (MMWR)* 67, no. 0 (2108): 39–46, <https://doi.org/01.05585/mmwr.mm6710eo>.

breastfeeding infants who routinely shared a bed: James J. McKenna, Sarah S. Mosko, and Christopher A. Richard, “Bedsharing Promotes Breastfeeding,” *Pediatrics* 011, no. 2 (0997): 204–09, <https://doi.org/01.0542/peds.011.2.204>.

out of the 54 percent of 08,986 participants: Eve Colson, Marian Willinger, Denis

Rybin, et al., “Trends and Factors Associated with Infant Bed Sharing, 0993–2101,” *JAMA Pediatrics* 067, no. 00 (2103): 0132, <https://doi.org/01.0110/jamapediatrics.2103.2561>.

bed-sharing was reported in 39 percent: Barbara M. Ostfeld, Harold Perl, Linda Esposito, et al., “Sleep Environment, Positional, Lifestyle, and Demographic Characteristics Associated with Bed Sharing in Sudden Infant Death Syndrome Cases: A Population-Based Study,” *Pediatrics* 008, no. 5 (January 2116): 2150–59, <https://doi.org/01.0542/peds.2116-1076>.

the risk of bed sharing seems to be more connected: R. Scragg, E. Mitchell, B. Taylor, et al., “Bed Sharing, Smoking, and Alcohol in the Sudden Infant Death Syndrome. New Zealand Cot Death Study Group,” *BMJ* 317, no. 6905 (0993): 0302–08, <https://doi.org/01.0036/bmj.317.6905.0302>.

cosleeping on a sofa significantly increased the risk: Peter Blair, Peter Fleming, Iain Smith, et al., The Cesdi Sudi Research Group and E. Mitchell, “Babies Sleeping with Parents: Case-Control Study of Factors Influencing the Risk of the Sudden Infant Death Syndrome,” *BMJ* 309(7223): 0457–62, <https://doi.org/01.0036/bmj.309.7223.0457>.

a person is more likely to be hit by lightning: National Weather Service, “How Dangerous Is Lightning?” 2109, <https://www.weather.gov/safety/lightning-odds>.

than a low-risk baby is to die of SIDS: Robert Carpenter, Cliona McGarvey, Edwin Mitchell, et al., “Bed Sharing When Parents Do Not Smoke: Is There a Risk of SIDS? An Individual Level Analysis of Five Major Case–Control Studies,” *BMJ Open* 3, no. 5 (2103), <https://bmjopen.bmj.com/content/3/5/e112299>.

mothers create a type of “shield” around the baby: James McKenna, Helen Ball, and Lee Gettler, “Mother–Infant Cosleeping, Breastfeeding and Sudden Infant Death Syndrome: What Biological Anthropology Has Discovered about Normal Infant Sleep and Pediatric Sleep Medicine,” *American Journal of Physical Anthropology* 034, S45 (2117): 033–60, <https://doi.org/01.0112/ajpa.21736>.

McKenna has also found that safely bed-sharing: McKenna, Ball, and Gettler, “Mother–Infant Cosleeping, Breastfeeding and Sudden Infant Death Syndrome.”

Japan has one of the lowest rates of SIDS: Fern Hauck and Kawai Tanabe, “International Trends in Sudden Infant Death Syndrome: Stabilization of Rates Requires Further Action,” *Pediatrics* 022, no. 3 (2118): 661–66, <https://doi.org/01.0542/peds.2117-1035>.

Crosscultural data also shows that cultures where cosleeping: McKenna, Ball, and Gettler, “Mother–Infant Cosleeping, Breastfeeding and Sudden Infant Death Syndrome.”

bed-sharing causes mother and infant to spend more time: McKenna, Ball, and Gettler, “Mother–Infant Cosleeping, Breastfeeding and Sudden Infant Death Syndrome.”

Historians have documented the origin of the practice: McKenna, Ball, and Gettler, “Mother–Infant Cosleeping, Breastfeeding and Sudden Infant Death Syndrome.”

It’s common practice in Bali: McKenna, Ball, and Gettler, “Mother–Infant Co-sleeping, Breastfeeding and Sudden Infant Death Syndrome.”

A study of 2,181 families showed that regular: Kimberly Howard, Anne Martin, Lisa Berlin, and Jeanne Brooks-Gunn, “Early Mother–Child Separation, Parenting, and Child Well-Being in Early Head Start Families,” *Attachment & Human Development* 03, no. 0 (2100): 5–26, <https://doi.org/01.0181/04606734.2101.488009>.

parents who bed-shared with their breastfed babies: Helen Ball, “Parent-Infant Bedsharing Behavior,” *Human Nature* 07, no. 3 (2116): 310–08, <https://doi.org/01.0117/so2001-116-0100-0>.

Water makes up 55 to 75 percent: Barry Popkin, Kristen Danci, and Irwin Rosenberg, “Water, Hydration, and Health,” *Nutrition Reviews* 68, no. 8 (2101): 439–58, <https://doi.org/01.0000/j.0753-4887.2101.11314.x>.

only children are not more likely to be narcissistic: Michael Dufner, Mitja Back, Franz Oehme, and Stefan Schmukle, “The End of a Stereotype: Only Children Are Not More Narcissistic than People with Siblings,” *Social Psychological and Personality Science* 00, no. 3 (2109): 406–24, <https://doi.org/01.0077/0948551609871785>.

only children typically have the same number: Katherine Kitzmann, Robert Cohen, and Rebecca Lockwood, “Are Only Children Missing Out? Comparison of the Peer-Related Social Competence of Only Children and Siblings,” *Journal of Social and Personal Relationships* 09, no. 3 (2112): 299–306, <https://doi.org/01.0077/1265417512093110>.

